

(NOT TO BE OPENED BEFORE TIME OR TILL ASKED TO DO SO)

PHDURS-EE-2013

SUBJECT : Chemistry

10013



Sr. No. ....

Time : 1½ Hours

Max. Marks : 100

Total Questions : 100

Candidate's Name \_\_\_\_\_ Date of Birth \_\_\_\_\_

Father's Name \_\_\_\_\_ Mother's Name \_\_\_\_\_

Roll No. (in figures) \_\_\_\_\_ (in words) \_\_\_\_\_


Date of Examination \_\_\_\_\_ Option Attempt (Under Part-II) \_\_\_\_\_

\_\_\_\_\_  
(Signature of the Candidate)\_\_\_\_\_  
(Signature of the Invigilator)

**CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.**

1. **Part-I** (Question No. 1 to 40) is compulsory. **Part-II** (Question Nos. 41 to 100) is Optional. From Part-II, the candidate is to attempt 60 questions from any **One Option** out of the **three Optional parts** i.e. either from Option "A" or "B" or "C". All questions carry equal marks.
2. All the candidates **must return** the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.
4. The candidate **must not** do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers **Must Not** be ticked in the question booklet.
5. **Only black or blue ball point pen is to be used in the OMR Answer-Sheet.**
6. For each correct answer, the candidate will get full credit. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer. There will be No Negative marking.
7. *Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet. Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.*

revised for  
jumble of  
21/9/2013




**PART – I**  
**(COMPULSORY)**

1. The geometry of  $IF_8^-$  ion is :
  - (1) Pyramidal
  - (2) Tetrahedral
  - (3) Trigonal bipyramidal
  - (4) Square antiprismatic
2. Which of the following statement is *false* ?
  - (1)  $[Cu(en)_2]^{2+}$  is more stable than  $[Cu(NH_3)_4]^{2+}$
  - (2)  $[FeF]^{2+}$  is stable than  $[FeCl]^{2+}$
  - (3)  $[Fe(CN)_6]^{4-}$  is less stable in comparison to  $[Fe(CN)_6]^{3-}$
  - (4)  $[Cu(NH_3)_4]^{2+}$  is less stable than  $[Cd(NH_3)_4]^{2+}$
3. The coordination numbers of Ti(N) and  $O^{2-}$  in rutile are, respectively :
  - (1) 6 and 3
  - (2) 3 and 6
  - (3) 2 and 4
  - (4) 4 and 2
4. Racemization of a chiral complex such as  $[Cr(ox)_3]^{3-}$  is least likely to occur by :
  - (1) a dissociative pathway
  - (2) a pathway involving a 5-coordinate species in which one  $ox^{2-}$  ligand is monodentate
  - (3) the Ray-Dutt twist mechanism
  - (4) the Bailar twist mechanism
5. In the base-catalysed substitution of  $Cl^-$  by  $OH^-$  in  $[Co(NH_3)_5Cl]^{2+}$  under strongly basic conditions, the first step in the mechanism is :
  - (1) conversion of an ammine to amido ligand
  - (2) substitution of  $Cl^-$  by  $[OH]^-$
  - (3) dissociation of  $Cl^-$  to give a 5-coordinate intermediate
  - (4) association of  $[OH]^-$  to give a 7-coordinate intermediate
6. In tetrahedral complexes, which orbital is involved in  $\sigma$  as well as  $\pi$  bond according to MO theory :
  - (1)  $e$
  - (2)  $t_2$
  - (3)  $a_1$
  - (4)  $b$

7. The term symbol for ground state of *Ni* is :
- (1)  ${}^7S_3$                       (2)  ${}^3F_4$                       (3)  ${}^3P_0$                       (4)  ${}^7F_2$
8. In which of the following configuration, the orbital contribution is quenched in octahedral field ?
- (1)  $t_{2g}^4 e_g^2$                       (2)  $t_{2g}^6 e_g^1$                       (3)  $t_{2g}^4$                       (4)  $t_{2g}^5 e_g^2$
9. Which of the following does *not* possess bridged CO ?
- (1)  $CO_2(CO)_8$                       (2)  $Fe_3(CO)_{12}$                       (3)  $Os_3(CO)_{12}$                       (4)  $Fe_2(CO)_9$
10. Which of the following will have highest CO stretching frequency ?
- (1)  $Cr(CO)_6$                       (2)  $Mn(CO)_6^+$                       (3)  $V(CO)_6^-$                       (4)  $Fe(CO)_4^{2-}$
11. At 25°C which of the following substance has the lowest molar entropy ?
- (1)  $N_2$  (gas)                      (2) Mg(s)                      (3)  $C_6H_6$ (l)                      (4)  $CCl_4$ (g)
12. The unit of the rate and rate constant are the same for a reaction of order :
- (1) 0                      (2) 1                      (3)  $\frac{1}{2}$                       (4) 2
13. Walden role is given by :
- (1) product of equivalent conductance and viscosity  
 (2) product of molarity and viscosity  
 (3) sum of viscosity and ionic conductance  
 (4) product of molarity and molecular mass
14. In Rice-Herzfeld mechanism of decomposition of acetaldehyde, the order of reaction is :
- (1) 1/2                      (2) 1                      (3) 3/2                      (4) 2
15. Clausius-Clapeyron equation is given by :
- (1)  $\log \frac{p_2}{p_1} = \frac{\Delta H_{vap}}{2.303 R} \left[ \frac{T_2 - T_1}{T_1 \times T_2} \right]$                       (2)  $\log \frac{p_1}{p_2} = \frac{\Delta H_{vap}}{2.303 R} \left[ \frac{T_1 - T_2}{T_1 \times T_2} \right]$
- (3)  $\log \frac{p_2}{p_1} = \frac{\Delta H_{vap}}{2.303} \left[ \frac{T_1 - T_2}{T_1 + T_2} \right]$                       (4)  $\log \frac{p_2}{p_1} = \frac{\Delta H_{vap}}{2.303} \left[ \frac{T_1 + T_2}{T_1 - T_2} \right]$



16. Which quantum number does not arise from solution of Schrodinger equation ?  
 (1) Principal quantum number (2) Spin quantum number  
 (3) Magnetic quantum number (4) Azimuthal quantum number
17. If length of the one dimensional box is halved, the energy of the particle will become :  
 (1) Half (2) Doubled  
 (3) Four times (4) One fourth
18. The degeneracy of energy level with energy equal to  $\frac{6h^2}{8na^2}$  is :  
 (1) 2 (2) 3 (3) 6 (4) 9
19. Unit of equivalent conductivity is :  
 (1)  $\text{ohm cm}^2 \text{eq}^{-1}$  (2)  $\text{ohm}^{-1} \text{cm}^2 \text{eq}^{-1}$   
 (3)  $\text{ohm cm}^{-2} \text{eq}^{-1}$  (4)  $\text{ohm}^{-1} \text{cm}^{-2} \text{eq}^{-1}$
20. If  $K$  = equilibrium constant,  $Q$  = reaction quotient and  $G$  = Gibb's free energy, which of the following is true for a spontaneous reaction ?  
 (1)  $\Delta G < \Delta G^\circ$  (2)  $\Delta G > \Delta G^\circ$  (3)  $K > Q$  (4)  $K < Q$
21. The two compounds shown below are :



- (1) diastereomers (2) enantiomers  
 (3) identical (4) conformational isomers
22. Which of the following is *not true* about enantiomers ? They have the same :  
 (1) Melting Point (2) Boiling Point  
 (3) Specific rotation (4) Density

23. When benzyl chloride is treated with ethanolic KCN, benzyl ethyl ether is produced along with benzyl cyanide. The most likely mechanism for the reaction would be :
- (1)  $SN^2$  (2)  $SN^1$   
(3)  $SN^i$  (4) Both  $SN^1$  and  $SN^2$
24. The carbene which adds stereo specifically to a double bond is in the ..... state.
- (1) singlet (2) doublet (3) triplet (4) free radical
25. The compound which would undergo nitration at ortho and para position with rates lower than that of benzene is :
- (1) Cinnamic acid (2) Toluene (3) Phenol (4) Benzoic acid
26. M-effect is :
- (1) Resonance effect (2) Inductive effect  
(3) No bond resonance (4) Electromeric effect
27. The type of linkage present in porcion is :
- (1) H-bonding (2) Covalent linkage  
(3) Ionic bond (4) Coordinate bond
28. Epimers differ in :
- (1) C - 1 (2) C - 2  
(3) C - 1 and C - 2 (4) None of the above
29. In the conversion of a Grignard reagent into an aldehyde the other component used is :
- (1) ethyl formate (2) ethyl acetate (3) ethyl cyanide (4)  $CO_2$
30. Sulphonation of benzene differs from most other electrophilic substitution reaction in that the reaction :
- (1) is reversible  
(2) occurs with explosive violence  
(3) requires elevated temperature  
(4) requires Lewis acid catalyst

31. The wave length of de-Broglie's wave associated with a moving proton of mass  $1.66 \times 10^{-27}$  kg and kinetic energy of  $5 \times 10^{-27}$  J is :
- (1)  $162.65 \times 10^{-8}$  m                      (2)  $16.265 \times 10^{-8}$  m  
(3) 16.265 m                                  (4) 1.6265 m
32. The pure rotational spectrum of gaseous HCl consists of a series of equally spaced lines separated by  $20.80 \text{ cm}^{-1}$ . The value of rotational constant is :
- (1)  $20.80 \text{ cm}^{-1}$       (2)  $10.40 \text{ cm}^{-1}$       (3)  $5.20 \text{ cm}^{-1}$       (4)  $2 \text{ cm}^{-1}$
33. Which of the following molecules has lowest vibrational stretching frequency ?
- (1)  $^1\text{H} \text{}^{35}\text{Cl}$       (2)  $^2\text{D} \text{}^{35}\text{Cl}$       (3)  $^1\text{H} \text{}^{36}\text{Cl}$       (4)  $^1\text{H} \text{}^{37}\text{Cl}$
34. The proton nmr spectrum of propane will consist of :
- (1) a triplet and a singlet                      (2) a triplet and a quartet  
(3) a doublet and a sextet                      (4) a triplet and a septet
35. To check that a secondary alcohol has been completely oxidized to a ketone you can :
- (1) check out the IR spectrum has absorptions at  $3500 \text{ cm}^{-1}$  and  $1650 \text{ cm}^{-1}$   
(2) check out the IR spectrum has no absorptions at  $3500 \text{ cm}^{-1}$  and  $1650 \text{ cm}^{-1}$   
(3) check out the IR spectrum has no absorptions at  $3500 \text{ cm}^{-1}$   
(4) check out the IR spectrum has no absorptions around  $1650 \text{ cm}^{-1}$
36. The  $\beta$ -isomer of hydrated trisglycinato cobalt (III) is ..... in colour consisting of two bands.
- (1) Red                      (2) Violet                      (3) Yellow                      (4) Blue
37. Which listed below gives only spin active nuclei ?
- (1)  $^1\text{H}, ^{13}\text{C}, ^{19}\text{F}$                                   (2)  $^2\text{H}, ^{12}\text{C}, ^{19}\text{F}$   
(3)  $^1\text{H}, ^2\text{H}, ^{12}\text{C}$                                   (4)  $^1\text{H}, ^{12}\text{C}, ^{19}\text{F}$

38. The position of the characteristic carbonyl stretching absorption bands in the IR spectrum of  and  are observed at :

- (1)  $1715\text{ cm}^{-1}$  and  $1680\text{ cm}^{-1}$                       (2)  $1680\text{ cm}^{-1}$  and  $1715\text{ cm}^{-1}$   
 (3)  $1740\text{ cm}^{-1}$  and  $1715\text{ cm}^{-1}$                       (4)  $1715\text{ cm}^{-1}$  and  $1740\text{ cm}^{-1}$

39. The lowest energy transition for tetrahedral complex of  $Mn^{2+}$  is :

- (1)  ${}^3A_2 \rightarrow T_1$               (2)  ${}^4T_1 \rightarrow {}^4A_2$               (3)  ${}^3A_2 \rightarrow {}^3T_2$               (4)  ${}^4A_2 \rightarrow {}^4T_2$

40. The cis isomers often have ..... molar absorptivity values for  $d \rightarrow d$  transitions than trans isomers.

- (1) Larger              (2) Smaller              (3) Equal              (4) None of the above

### PART - II

#### (OPTIONAL)

#### OPTION - A : INORGANIC CHEMISTRY

41. In  $CO_2$  molecule, the band at  $1340\text{ cm}^{-1}$  due to fermiresonance, has band maxima (doublet) at :
- (1)  $1286$  and  $1388\text{ cm}^{-1}$                       (2)  $1276$  and  $1398\text{ cm}^{-1}$   
 (3)  $1277$  and  $1397\text{ cm}^{-1}$                       (4) None of these
42. In  $AB_5$  type TBP molecules, the number of IR active stretching vibrations are :
- (1) Three              (2) Four              (3) Two              (4) Five
43. In thiocynato complexes, the  $C \equiv N$  stretching frequencies are ..... than in isothiocynato complexes.
- (1) Higher              (2) Lower              (3) Similar              (4) None of these
44. Value of 'g' for an atom having ground state term symbol  ${}^2P_{3/2}$  will be :
- (1) 2.0              (2) 1.33              (3) 1.73              (4) 2.25
45. In EPR spectrum of bis (salicyladimine) copper (II), the hyperfine structure of each major peak consists of :
- (1) Nine subpeaks                      (2) Fifteen subpeaks  
 (3) Eleven subpeaks                      (4) Ten subpeaks



46. Quadrupole splitting is *not* observed in the MB spectrum of :  
(1)  $Fe(CO)_5$       (2)  $FeSO_4$       (3)  $K_3[Fe(CN)_6]$       (4)  $FeCl_3$
47. The radical anion  $[ON(SO_3)_2]^{2-}$  shows in ESR :  
(1) A triplet hyperfine structure from nitrogen  
(2) Hyperfine splitting of 13.05 gauss  
(3) No splitting due to S and O  
(4) All of the above
48. Which is *correct* order of chemical shift ( $\delta$ ) decrease in MB spectra ?  
(1)  $Cl^- > O^{2-} > N^{3-} > CN^-$       (2)  $CN^- > O^{2-} > N^{3-} > Cl^-$   
(3)  $Cl^- > CN^- > O^{2-} > N^{3-}$       (4)  $CN^- > N^{3-} > O^{2-} > Cl^-$
49. Which does *not* apply to mass spectrometry ?  
(1) Magnetic field      (2) Acceleration potential  
(3) Microwaves      (4) Ionization and fragmentation
50. Which change is *not* detected by DTA ?  
(1) Polymer softening      (2) Desorption  
(3) Sublimation      (4) Loss of moisture
51. Stability of nucleus is due to :  
(1) Long-range forces      (2) Short-range forces  
(3) Pion cloud only      (4) None of the above
52. Which nuclear model can best explain that all elements with atomic number greater than 92 are radioactive ?  
(1) Liquid Drop Model      (2) Shell Model  
(3) Collective Model      (4) All of these
53. What is the total binding energy of  ${}^6_3Li$  nucleus having atomic mass 6.0170 amu ?  
(Mass of proton = 1.00727 a.m.u. and mass of neutron = 1.008665 amu)  
(1) 28.82 MeV      (2) 27.89 MeV      (3) 28.69 MeV      (4) 27.69 MeV

54. Spallation reactions are initiated by high speed :
- (1) Protons (2)  $\alpha$ -particles  
(3) Both Protons and  $\alpha$ -particles (4) None of these
55.  ${}_{13}^{27}\text{Al}$  is a stable Isotope. It is expected to disintegrate by :
- (1)  $\alpha$  - emission (2)  $\beta^-$  emission (3)  $\beta^+$  emission (4) Proton emission
56. The radioactivity detector based on light emission is :
- (1) Cloud Chamber (2) Ionization Chamber  
(3) Scintillation Counter (4) Solid State Detector
57. To which element, Neutron Activation Analysis is applicable ?
- (1) Magnesium (2) Niobium (3) Vanadium (4) Copper
58. The sensitivity of NAA depends upon :
- (1) Atomic cross section of particles (2) Flux of particles  
(3) Half life of Nuclide (4) All of these
59. Ionization Chamber uses lower operating voltage than :
- (1) Proportional Counters (2) Solid ion Chamber  
(3) Scintillation Counter (4) All of these
60. Which of the following Nuclei is *not* doubly magic ?
- (1)  ${}^4_2\text{He}$  (2)  ${}^{16}_8\text{O}$  (3)  ${}^{208}_{82}\text{Pb}$  (4)  ${}^{238}_{92}\text{U}$
61. The metal species present in Nitrogenase is :
- (1) Zinc (2) Molybdenum (3) Tungsten (4) Lead
62. Which of the following is used in Psychotropic drugs ?
- (1) Sodium fluoride (2) Lithium carbonate  
(3) Barium sulphide (4) Zinc oxide
63. CYTOCHROM P-450 enzyme contains metal :
- (1) Zinc (2) Copper (3) Cobalt (4) Iron

64. Deficiency of  $Zn$  causes the disease :
- (1) Convulsions
  - (2) Liver necrosis
  - (3) Dwarfism
  - (4) Kinky-hair syndrome
65. Oxymyoglobin contains :
- (1) Oxygen in hole of Porphyrin
  - (2) Oxygen bonded to  $Mg$
  - (3) Oxygen at trans position to histidine chain
  - (4) Oxygen not present at all
66. Photochemical Smog is caused by :
- (1) Oxides of Nitrogen
  - (2) Hydrocarbons
  - (3) Carbon monoxide
  - (4) Oxides of N, Hydrocarbons and  $CO$
67. Ozone depletion in Antarctica is due to the formation of :
- (1) Acrolin
  - (2) Peroxyacetylnitrate
  - (3)  $SO_2$  and  $SO_3$
  - (4) Chlorine nitrate
68. Silicosis is caused by :
- (1) Acid rain
  - (2) Depletion of Ozone
  - (3) Inhalation of aerosols
  - (4) Inhalation of  $SO_2$
69. Catechol type siderophore is :
- (1) Ferrichrome
  - (2) Enterobactin
  - (3) Ferrioxamine
  - (4) None of these
70. In the resting state, the level of  $Ca^{2+}$  near the muscle fibre is :
- (1) Very low
  - (2) Very high
  - (3) Medium
  - (4) No change
71. The heptacity of tropylium ion is :
- (1)  $n^5$
  - (2)  $n^1$
  - (3)  $n^7$
  - (4)  $n^3$
72. Fluxional behaviour in a molecule can be detected by :
- (1) IR spectroscopy
  - (2) X-rays
  - (3) NMR spectroscopy
  - (4) UV-Vis spectroscopy

73. Ziegler-Natta catalyst is :

- (1)  $TiCl_4 - AlEt_3$  (2)  $RhCl(PPh_3)_3$   
 (3)  $CO_2(CO)_8$  (4)  $PdCl_4^{2-}$

74. Electrophilic Carbene ligands are also called :

- (1) Fischer Carbene (2) Schrock Carbene  
 (3) Homonuclear Carbene (4) Heteronuclear Carbene

75. In Ferrocene, which metal orbital interacts with the composite ring orbitals  $C_pE_{1g}$  of ligand for the formation of covalent bonds :

- (1)  $4pz, 4px$  (2)  $3dxz, 3dyz$   
 (3)  $3dxy, 3d_{x^2-y^2}$  (4)  $(DS)_x, (DS)_y$

76. The C = C infrared absorption peak of  $[Mn(\eta^3-C_3H_5)(CO)_4]$  appears at :

- (1)  $1620\text{ cm}^{-1}$  (2)  $1570\text{ cm}^{-1}$  (3)  $1505\text{ cm}^{-1}$  (4)  $1520\text{ cm}^{-1}$

77. Which metal alkyne complex is  $4e^-$  donor ?

- (1)  $Pt^{II}Cl_2$  (p-toluidine)  $Bu^+C \equiv C Bu^+$   
 (2)  $Pt^0(PPh_3)_2(PhC \equiv CPh)$   
 (3)  $[C_2H_2CO_2(CO)_6]$   
 (4) None of the above

78. Transition metal alkene complexes are readily attacked by :

- (1) Electrophile  
 (2) Nucleophile  
 (3) Both Electrophile and Nucleophile  
 (4) No reaction with electrophile & nucleophile

79. Which of the following does *not* obey EAN rule ?

- (1)  $V(CO)_3(\pi-C_5H_5)(R_2C=C R_2)_2$  (2)  $Co(CO)_2(\pi-C_5H_5)$   
 (3)  $Fe(\sigma-C_5H_5)(\pi-C_5H_5)(CO)_3$  (4)  $Cr(C_6H_6)(CO)_3$

80. Proton NMR spectrum of  $(n^1Cp)(n^5Cp)Fe(CO)_2$  at ambient temperature shows :
- (1) Two singlets of almost equal intensity
  - (2) A singlet and a multiplet of equal intensity
  - (3) One singlet of high intensity
  - (4) Two multiplets of equal intensity
81. The current due to supporting electrolyte is called :
- (1) Residual Current
  - (2) Diffusion Current
  - (3) Migration Current
  - (4) Alternate Current
82. In anodic stripping voltametry, the concentration of metal ions is in the range of :
- (1)  $10^{-3}$  to  $10^{-6}$  m
  - (2)  $10^{-4}$  to  $10^{-7}$  m
  - (3)  $10^{-5}$  to  $10^{-8}$  m
  - (4)  $10^{-5}$  to  $10^{-10}$  m
83. The half wave potential for  $Cu^{2+}$  in 1 M NaOH is :
- (1) -1.12 V
  - (2) -0.41 V
  - (3) -1.53 V
  - (4) -1.46 V
84. The diffusion current in polarography is given by :
- (1)  $i_d = i_l - i_r$
  - (2)  $i_d = i_l + i_r$
  - (3)  $i_d = 2i_l - i_r$
  - (4)  $i_d = i_l - 2i_r$
85. Ion-selective membrane used in ion selective electrodes are :
- (1) Glass membranes
  - (2) Crystalline membranes
  - (3) Ion exchange resin membranes
  - (4) All of the above
86. The detection limit for anodic stripping voltametry is :
- (1)  $10^{-2}$  to  $10^{-4}$  m
  - (2)  $10^{-4}$  to  $10^{-6}$  m
  - (3)  $10^{-9}$  to  $10^{-10}$  m
  - (4)  $10^{-5}$  to  $10^{-7}$  m
87. Which of the following can be used as end point detection technique in Coulometric titrations ?
- (1) Potentiometry
  - (2) Amperometry
  - (3) Conductometry
  - (4) Potentio, ampero and conductometry
88. A rotating Pt electrode is preferred over DME in the titration involving :
- (1) Bromine
  - (2)  $Ag^+$  ion
  - (3)  $Fe^{2+}$  ion
  - (4) Br,  $Ag^+$  and  $Fe^{2+}$  all

89. In nuclear medicine imaging, radiopharmaceuticals are taken :  
(1) Intravenously (2) Orally  
(3) Both I and II (4) Neither I nor II
90. The mode of decay in radio Iodine-131 is :  
(1)  $\alpha$ -decay (2)  $\beta$ -decay (3)  $\gamma$ -decay (4) Neutron decay
91. The increased concentration of  $K^+$  in extra cellular fluid causes :  
(1) Hypokalemia (2) Hyperkalemia  
(3) Addison's disease (4) Dyspnea
92. Liver necrosis disease is caused by deficiency of :  
(1) Calcium (2) Chromium (3) Selenium (4) Cobalt
93. Cancer causing chemicals are :  
(1) Oxines & Azo compounds (2) Urethanes & nitrosoamines  
(3) Alkylating agents (4) All of these
94. Which iron salt has minimum interference with tetracyclin drug absorption in gut ?  
(1) Ferrous sulphate (2) Ferrous fumerate  
(3) Ferrous succinate (4) Ferric – EDTA
95. Which of the following is a antiviral drug ?  
(1) 1-methyl-2-mercaptoimidazole  
(2) 1-methylisatin-3-thiosemicarbazone  
(3) 2-formyl-pyridine thiosemicarbazone  
(4) Aspirin
96. 1-methyl-2-mercaptoimidazole is used as potential agent for :  
(1) Anti thyroid activity (2) Anti cancer activity  
(3) Anti bacterial activity (4) Anti malarial activity
97. The concentration of Lithium in Plasma should be :  
(1) 2.0 m mol/litre (2) 0.4 – 1.6 m mol/litre  
(3) 0.6 – 1.2 m mol/litre (4) 2.0 – 2.4 m mol/litre

98. Chemical name of Vitamin  $B_{12}$  is :
- (1) Cyanocobalamin
  - (2) Hydroxycobalamin
  - (3) Methylcobalamin
  - (4) Cyano-, hydroxy - and methyl cobalamin
99. Recommended Dietary allowances for a male (19-70 years) for Vitamin C is :
- (1) 15 mg
  - (2) 90 mg
  - (3) 20 mg
  - (4) 5.0 mg
100. Source for polyphenolic antioxidants are food such as :
- (1) fresh fruits and vegetables
  - (2) whole wheat cereals and tea
  - (3) vegetable oils
  - (4) eggs

**OPTION – B : PHYSICAL CHEMISTRY**

41. The radius of  ${}_{13}^{27}\text{Al}$  nucleus is :
- (1)  $4.5 \times 10^{-15}$  m
  - (2)  $4.5 \times 10^{-14}$  m
  - (3)  $4.5 \times 10^{-16}$  m
  - (4)  $4.5 \times 10^{-13}$  m
42. Let a molecule  $AB_6$  belongs to  $O_h$  point group. The point group that result if it is changed to  $AB_5C$  would be :
- (1)  $D_{4h}$
  - (2)  $C_{2v}$
  - (3)  $C_{4v}$
  - (4) None of these
43. Water molecule belongs to point group :
- (1)  $C_{3v}$
  - (2)  $C_{2v}$
  - (3)  $D_{4h}$
  - (4)  $D_{2h}$
44. The hyperfine splitting constant, 'a' utilized to map the molecular orbital occupied by unpaired electron is related by :
- (1)  $a = Q^e$
  - (2)  $A = Q/e$
  - (3)  $a = Q + e$
  - (4)  $A = Q - e$
- where  $e$  is the unpaired density on a carbon atom.
45. The Mössbauer spectra of  $K_4Fe(CN)_6$  and  $[K_3Fe(CN)_5NO]$  consist of, respectively :
- (1) one line each
  - (2) two lines each
  - (3) two and four lines
  - (4) one and two lines

46. The power output of a laser in which a 2.0 J pulse can be delivered in one nanosecond is :
- (1) 2.0 GW            (2) 0.20 GW            (3) 20.0 GW            (4) 0.02 GW
47. The Miller indices of crystal plane which cut through the crystal axis at  $(2a, 3b, c)$  are :
- (1) (236)            (2) (326)            (3) (623)            (4) (362)
48. Frenkel defects appear in crystals, in which :
- (1) positive ions are much larger than the negative ions  
(2) positive ions are equal to negative ions in size  
(3) negative ions are much larger than the positive ions  
(4) None of the above
49. The coordination number of an atom in a face-centred cubic unit cell is :
- (1) 1            (2) 6            (3) 8            (4) 12
50. The probability factor in collision theory of reaction rates should be interpreted in terms of :
- (1) Enthalpy            (2) Free energy            (3) Entropy            (4) Viscosity
51. NMR transition is shifted from the reference in a 400 MHz spectrometer by 529 Hz. The chemical shift is :
- (1) 1.32            (2) 5.29            (3) 1.82            (4) 7.58
52. If activation energy of a certain reaction is zero, then rate constant will be equal to :
- (1) infinity            (2)  $A$             (3) zero            (4)  $A^{-1}$
- where  $A$  is the frequency factor.
53. Which one of the following statements about ionization in mass spectrometer is *incorrect* ?
- (1) Gaseous atoms are ionized by bombarding them with high energy electrons  
(2) Atoms are ionized so they can be accelerated  
(3) Atoms are ionized so they can be deflected  
(4) It doesn't matter how much energy you use to ionize atoms
54. The region of an infra-red spectrum where many absorptions takes place is known as :
- (1) Thumb print region            (2) Hand print region  
(3) Finger print region            (4) Foot print region



55. No diffraction would result, if :  
(1)  $\lambda \ll 2d$       (2)  $\lambda \gg 2d$       (3)  $\lambda \approx 2d$       (4)  $\lambda \ll d$
56. Select the correction equation from the following :  
(1)  $\left(\frac{\partial V}{\partial T}\right)_S = \frac{C_v}{T} \left(\frac{\partial T}{\partial P}\right)_V$       (2)  $\left(\frac{\partial S}{\partial V}\right)_T = \frac{C_p}{T} \left(\frac{\partial T}{\partial V}\right)_P$   
(3)  $\left(\frac{\partial V}{\partial T}\right)_V = \frac{C_v}{T} \left(\frac{\partial T}{\partial P}\right)_V$       (4)  $\left(\frac{\partial S}{\partial V}\right)_P = \frac{C_p}{T} \left(\frac{\partial T}{\partial V}\right)_P$
57. Which of the following is *not* a state function ?  
(1) Work      (2) Heat      (3) Enthalpy      (4) Entropy
58. The fundamental vibrational frequency of a molecule is  $1035 \text{ cm}^{-1}$ . Its force constant would be :  
(1)  $4\pi^2 c \mu^2 (1035) \times 10^4$       (2)  $4\pi^2 c^2 \mu^2 (1035)^2 \times 10^2$   
(3)  $4\pi^2 c^2 \mu (1035)^2 \times 10^4$       (4)  $4\pi^2 c^2 \mu (1035)^2 \times 10^2$
59. The pH of a solution is 6. Acid is added to decrease the pH to 4. The increase in hydrogen ion concentration is :  
(1) Hundred times      (2) Two times  
(3) Thousand times      (4) Ten times
60. The quantum yield of photochemical gas reaction  $2 \text{ HI} \rightleftharpoons \text{H}_2 + \text{I}_2$  at wavelength  $2400 \text{ \AA}$  is :  
(1) 0.20      (2)  $10^3$       (3) 10      (4) 2
61. Which of the following statement is *correct* ?  
(1) A triple point is invariant  
(2) A triple point is monovariant  
(3) A triple point is also called incongruent melting point  
(4) Eutectic point is same as triple point
62. Mean free path of a gas molecule is :  
(1) independent of pressure  
(2) inversely proportional to temperature  
(3) directly proportional to pressure  
(4) None of these

63. Van't Hoff equation ; (at  $c \rightarrow 0$ ) for predicting molar mass of a polymer solution reduces to :

$$(1) \lim_{c \rightarrow 0} \left( \frac{\pi}{c} \right) = \frac{R}{M}$$

$$(2) \lim_{c \rightarrow 0} \left( \frac{\pi}{c} \right) = \frac{T}{M}$$

$$(3) \lim_{c \rightarrow 0} \left( \frac{\pi}{c} \right) = \frac{RT}{M}$$

$$(4) \lim_{c \rightarrow 0} \left( \frac{\pi}{c} \right) = \frac{RM}{T}$$

Where  $\pi$  is the osmotic pressure.

64. The heterogeneity of the polymer sample is called its :

(1) Polydispersity index

(2) Monodispersity

(3) Average molecular mass

(4) Polydispersity

65. Oriental polarizability  $\alpha$ , is related to temperature  $T$ , as :

$$(1) \alpha = \frac{\mu}{3kT}$$

$$(2) \alpha = \frac{\mu^2}{3kT}$$

$$(3) \alpha = \frac{\mu}{kT}$$

$$(4) \alpha = \mu kT$$

where all the symbols have usual significance.

66.  $\psi_{21(-1)}$  represents :

(1)  $2s$  orbital

(2)  $2p_x$  orbital

(3)  $2p_y$  orbital

(4)  $2p_z$  orbital

67. The average of a measurable property  $p_x$ , can be determined by employing relation :

$$(1) \langle p_x \rangle = \frac{\int \hat{p}_x \phi \phi^* d\tau}{\int \phi \phi^* d\tau}$$

$$(2) \langle p_x \rangle = \frac{\int \phi \hat{p}_x \phi^* d\tau}{\int \phi \phi^* d\tau}$$

$$(3) \langle \hat{p}_x \rangle = \frac{\int \phi \phi^* \hat{p}_x d\tau}{\int \phi \phi^* d\tau}$$

(4) None of the above

68.  $\left[ x, \frac{d}{dx} \right]$  will yield :

(1) zero

(2) 1

(3) -1

(4) 2

69. The Hamiltonian operator for a Helium atom is expressed by :

$$(1) \hat{H} = \frac{-h^2}{2m} (\nabla_1^2 + \nabla_2^2) + \frac{e^2}{r_{12}}$$

$$(2) \hat{H} = \frac{-h^2}{2m} (\nabla_1^2 + \nabla_2^2) + \frac{ze^2}{r_1} + \frac{ze^2}{r_2}$$

$$(3) \hat{H} = \frac{-h^2}{2m} (\nabla_1^2 + \nabla_2^2) - \frac{ze^2}{r_1} - \frac{ze^2}{r_2} + \frac{e^2}{r_{12}}$$

$$(4) \hat{H} = \frac{-h^2}{2m} (\nabla_1^2 + \nabla_2^2) - \frac{e^2}{r_{12}} + \frac{ze^2}{r_1} + \frac{ze^2}{r_2}$$

where  $\nabla_1$  and  $\nabla_2$  are Laplacien operators for electrons 1 and 2 respectively. All other symbols have usual significance.

70. The Eigen value is/can :

- |                     |                                     |
|---------------------|-------------------------------------|
| (1) always positive | (2) always negative                 |
| (3) be zero         | (4) be positive as well as negative |

71. Marcus refined the RRK theory by taking into consideration :

- (1) vibrations of the energized molecule
- (2) rotations of the energized molecule
- (3) all vibrations and rotation of the energized molecule
- (4) None of these

which in turn led to RRKM theory

72. The steric factor, P is related to Entropy of activation,  $\Delta S^\ddagger$  by :

$$(1) P = \frac{RT}{ZNh} \cdot e^{\Delta S^\ddagger/R}$$

$$(2) P = \frac{RT}{ZNh} \cdot e^{-\Delta S^\ddagger/R}$$

$$(3) P = \frac{RT}{h} \cdot e^{\Delta S^\ddagger/RT}$$

$$(4) P = \frac{R}{ZNh} \cdot e^{\Delta S^\ddagger/RT}$$

73. The Gibbs adsorption equation is :

$$(1) \Gamma = \frac{-RT}{C} \cdot \frac{dc}{dr}$$

$$(2) \Gamma = \frac{-CT}{R} \cdot \frac{dr}{dc}$$

$$(3) \Gamma = \frac{-C}{RT} \cdot \frac{dr}{dc}$$

$$(4) \Gamma = \frac{-CT}{R} \cdot \frac{dc}{dr}$$

where all the notations have usual significance.

74. The cell potential is a :

(1) Thermodynamic property

(2) Colligative property

(3) Extensive property

(4) Intensive property

75. How many normal modes of vibration are possible for benzene molecule ?

(1) 6

(2) 30

(3) 12

(4) 8

76. Synthetic fibres like nylon-66 are very strong because :

(1) They have linear molecules consisting of very long chains

(2) They have high molecular weights and high melting points

(3) They have a high degree of cross-linking by strong carbon-carbon bond

(4) They have linear molecules interlinked with forces like hydrogen bonding

77. Polyethene is :

(1) Thermosetting

(2) Thermoplastic

(3) Both (1) and (2)

(4) None of these

78. A solid acts as an adsorbent because it has :

(1) a definite shape

(2) a high lattice energy

(3) unsaturated valencies

(4) small pores in it

79. According to Langmuir adsorption isotherm, the amount of gas adsorbed at very high pressure :

(1) goes on decreasing with pressure

(2) goes on increasing with pressure

(3) increases first and decreases later with pressure

(4) reaches a constant limiting value

80. Lyophilic sols are more stable than lyophobic sols because :
- (1) The colloidal particles are solvated
  - (2) The colloidal particles have positive charge
  - (3) The colloidal particles have no charge
  - (4) There are strong electrostatic repulsions between the negatively charged colloidal particles
81. Which statement corresponds to the case where the chemical shift difference between two coupling protons is less than five times the coupling constant ?
- (1) An Ax pattern is observed
  - (2) An AB pattern is observed
  - (3) A first order spectrum is expected
  - (4) An undistorted binomial pattern is expected
82. Which of the following statement is false about NMR experiment ?
- (1) The energy difference between two spin states depends on the strength of magnetic field
  - (2) When energy absorption occurs, the nuclei are said to be in resonance with the electromagnetic radiation
  - (3) The energy required to flip the spin of a proton is in the infrared region of the electromagnetic radiation
  - (4) None of these
83. The number of microstates for distributing three atoms among energy states, having three quanta of energy are :
- (1) 1
  - (2) 6
  - (3) 10
  - (4) 3
84. The rotational energy possessed by atom having one degree of atom is :
- (1)  $RT$
  - (2)  $kT$
  - (3)  $\frac{1}{2}RT$
  - (4)  $\frac{1}{2}kT$
85. Translational partition function,  $q_t$  is expressed by :
- (1)  $q_t = \frac{(2\pi mkT)^{3/2} V}{RT}$
  - (2)  $q_t = \frac{(2\pi mkT)^{3/2}}{RT}$
  - (3)  $q_t = \frac{(2\pi mRT)^{3/2} V}{RT}$
  - (4)  $q_t = \frac{(2\pi mRT)^{3/2}}{T}$

86. The step down ladder operator is :

$$(1) \hat{J}_+ = \hat{J}_x + \hat{J}_y \quad (2) \hat{J}_+ = \hat{J}_x - \hat{J}_y \quad (3) \hat{J}_- = \hat{J}_x + i\hat{J}_y \quad (4) \hat{J}_- = \hat{J}_x - i\hat{J}_y$$

where all the symbols have usual significance.

87. Molecules orbital theory :

- (1) underestimates the importance of covalent structures
- (2) overestimates the importance of ionic structures
- (3) puts equal importance on both ionic and covalent structures
- (4) None of the above

88. Operators  $\hat{A}$  and  $\hat{B}$  are said to be commutative, if :

$$(1) \hat{A} + \hat{B} = 0 \quad (2) \hat{A} - \hat{B} = 0$$

$$(3) \hat{A} \hat{B} + \hat{B} \hat{A} = 0 \quad (4) \hat{A} \hat{B} - \hat{B} \hat{A} = 0$$

89. Which of the following is *correct* ?

$$(1) [\hat{L}^2, \hat{L}_z] > 0 \quad (2) [\hat{L}^2, \hat{L}_z] < 0$$

$$(3) [\hat{L}^2, \hat{L}_z] = 0 \quad (4) [\hat{L}^2, \hat{L}_z] = i\hbar \hat{L}_x$$

90. Which of the following is *true* ?

$$(1) q_{tr} \gg q_{rot} \gg q_{vib} \gg q_{elect} \quad (2) q_{tr} \gg q_{vib} > q_{rot} > q_{elect}$$

$$(3) q_{tr} \ll q_{rot} \ll q_{vib} \ll q_{elect} \quad (4) q_{tr} < q_{vib} < q_{rot} < q_{elect}$$

where  $q_{tr}$ ,  $q_{rot}$ ,  $q_{vib}$  and  $q_{elect}$  are translational, rotational, vibrational and electronic partition function.

91. When *Pt* and *Co* are electrically connected, which one gets corroded ?

- (1) *Pt*
- (2) *Co*
- (3) Cannot decide
- (4) None

92. Pipes of different materials, such as copper and steels, should not be embedded in a trench in close proximity to avoid :

- (1) deposition of copper on steel pipe
- (2) depassivation of steel
- (3) corrosion of copper pipes
- (4) galvanic corrosion

93. If moisture and dirt entrapment is a major problem, it would be good practice to :  
(1) Spot weld      (2) Skip weld      (3) Stitch weld      (4) Butt weld
94. The number of  $\alpha$  and  $\beta$  particles emitted by  ${}_{81}^{218}\text{Ra}$  in changing to a stable isotope of  ${}_{82}^{206}\text{Pb}$  will be :  
(1) 3 and 2      (2) 2 and 4      (3) 3 and 4      (4) 3 and 1
95. Milk is a/an :  
(1) Emulsion      (2) Gel      (3) Suspension      (4) Pure solution
96. At temperature near absolute zero, gaseous particles possess only :  
(1) Translational energy      (2) Vibrational energy  
(3) Rotational energy      (4) Rotational and vibrational energy
97. Lattice strength of various types of crystals vary as :  
(1) Ionic > covalent > metallic > molecular  
(2) Covalent > metallic > ionic > molecular  
(3) Metallic > covalent > ionic > molecular  
(4) Covalent > ionic > metallic > molecular
98. The energy per mole of light having wavelength of 85 nm is :  
(1)  $1.207 \times 10^6 \text{ J mole}^{-1}$       (2)  $1.307 \times 10^6 \text{ J mole}^{-1}$   
(3)  $1.407 \times 10^6 \text{ J mole}^{-1}$       (4)  $1.507 \times 10^6 \text{ J mole}^{-1}$
99. Which of the following has been used in the manufacture of non-inflammable photographic films ?  
(1) Cellulose nitrate      (2) Cellulose xanthate  
(3) Cellulose perchlorate      (4) Cellulose acetate
100. Which of the following is an irreversible cell ?  
(1)  $\text{Zn} / \text{Zn}^{2+} / \text{AgCl} / \text{Ag}$       (2)  $\text{Zn} / \text{H}_2\text{SO}_4 / \text{Ag}$   
(3)  $\text{Zn} / \text{Zn}^{2+} // \text{Cd}^{2+} / \text{Cd}$       (4)  $\text{Cd} / \text{Cd}^{2+} // \text{KCl}, \text{Hg}_2\text{Cl}_2(\text{s}) / \text{Hg}$

**OPTION – C : ORGANIC CHEMISTRY**

41. Which is *not* an anticancer drug ?  
 (1) Vincristine (2) Cyclophosphamide  
 (3) Doxorubicin (4) Gabapentin
42. What kind of spectroscopy is FT NMR ?  
 (1) Absorption (2) Emission (3) Both of these (4) None
43. The presence of a bromine is indicated in a compound if its mass spectrum shows M and M + 2 peaks in the intensity ratio :  
 (1) 2 : 1 (2) 3 : 1 (3) 1 : 1 (4) 1 : 2
44. LAH in combination with  $AlCl_3$  can be used to convert diarylketone ( $Ar_2CO$ ) into :  
 (1)  $Ar_2CHOH$  (2)  $Ar_2CH_2$  (3)  $ArCHOAr$  (4)  $Ar - Ar$
45. 1,3-Dithiane is a structural equivalent of :  
 (1) Acylcarbanion (2) Formylcarbanion  
 (3) Acyl carbonium ion (4) Formylcarbonium ion
46. Select the right decreasing order of nucleophilicity :  
 (1)  $CH_3 - \overset{\ominus}{C}H_2 > \overset{\ominus}{N}H_2 > CH \equiv \overset{\ominus}{C} > \overset{\ominus}{O}H$   
 (2)  $CH \equiv \overset{\ominus}{C} > \overset{\ominus}{N}H_2 > CH \equiv \overset{\ominus}{C} > \overset{\ominus}{O}H$   
 (3)  $\overset{\ominus}{O}H > \overset{\ominus}{N}H_2 > CH \equiv \overset{\ominus}{C} > CH_3 - \overset{\ominus}{C}H_2$   
 (4)  $\overset{\ominus}{N}H_2 > CH \equiv \overset{\ominus}{C} > \overset{\ominus}{O}H > CH_3 - \overset{\ominus}{C}H_2$
47. The ratio M | M + 2 for the presence of chlorine in a compound in its mass spectrum is :  
 (1) 3 : 1 (2) 1 : 2 (3) 4 : 2 (4) 2 : 1
48. Which is right about stretching frequencies of C = C and C = O in i. r. spectroscopy from intensity point of view ?  
 (1)  $V_{C=O}$  is stronger than  $V_{C=C}$   
 (2)  $V_{C=O}$  is weaker than  $V_{C=C}$   
 (3)  $V_{C=O}$  and  $V_{C=C}$  have equal intensity  
 (4) None of these



49. What is the decreasing order of chemical shifts for protons among these compounds ?  
(1) Alkynes > Alkanes > Alkenes      (2) Alkynes > Alkenes > Alkanes  
(3) Alkanes > Alkynes > Alkenes      (4) Alkenes > Alkynes > Alkanes
50. Mass spectroscopy requires a minimum sample size of :  
(1) Micrograms      (2) Nanograms      (3) Picograms      (4) Grams
51. Internal reference for phosphorus-31 is :  
(1)  $H_3PO_2$  (85%)      (2)  $H_3PO_4$  (85%)      (3)  $H_3PO_3$  (85%)      (4) None of these
52. Oct-4-ene shows C = C frequency in its i. r. spectrum at :  
(1)  $1680 - 1600 \text{ cm}^{-1}$  (vw)      (2)  $1680 - 1600 \text{ cm}^{-1}$  (s)  
(3)  $1680 - 1600 \text{ cm}^{-1}$  (m)      (4) No peak in this region
53. Continuous wave NMR spectroscopy involves :  
(1) simultaneous detection of all resonances  
(2) sequential detection of resonances of nuclei  
(3) first simultaneous followed by sequential detection of resonances  
(4) sometimes sequential and sometimes simultaneous detection of resonances
54. The  $C_{60}$  fullerene shows lesser number of peaks in the i. r. spectrum because :  
(1) It contains a graphite like structure  
(2) It is asymmetric  
(3) It contains  $sp^3$ ,  $sp^2$  and  $sp$  carbons  
(4) It has a symmetrical structure
55. Carbonyl compounds exhibit the transition :  
(1)  $\sigma - \sigma^*$   $2 \pi - \pi^*$       (2)  $\sigma - \pi^*$ ,  $\pi - \pi^*$ ,  $n - \pi^*$   
(3)  $\sigma - \sigma^*$ ,  $n - \sigma^*$ ,  $\pi - \pi^*$       (4) None of these
56. By which of these, acetophenone can be converted to phenol :  
(1) m-CPBA followed by base catalyzed hydrolysis  
(2) Conc.  $HNO_3$   
(3) Iodine and  $NaOH$   
(4) singlet oxygen followed by base catalyzed hydrolysis

57. Diazomethane with acetylene gives :  
(1) Pyrazole      (2) Pyrazoline      (3) Piperidine      (4) Pyrimidine
58. Which is used for treating Gout ?  
(1) Reserpine      (2) Atropine      (3) Colchine      (4) None
59. Cinnamoyl alcohol upon treatment with lead tetraacetate gives :  
(1) Acetophenone      (2) Cinnamic acid  
(3) Propanal      (4) Cinnamaldehyde
60. Which is a strong base ?  
(1) Aniline      (2) Cyclohexylamine  
(3) Pyrrole      (4) Quinoline
61. In  $SN^2$  displacement on methyl bromide, which is most effective ?  
(1)  $C_2H_5O^\ominus$       (2)  $HO^\ominus$       (3)  $C_6H_5O^\ominus$       (4)  $CH_3COO^\ominus$
62. Of these which reacts fastest with N-bromosuccinimide (NBS) ?  
(1) Toluene      (2) Methane      (3) Pyridine      (4) Benzene
63. Generally, an increase in solvent polarity for the reaction between alkylhalide and  $OH^\ominus$  :  
(1) Increases the rate of  $SN^1$  reaction  
(2) Decreases the rate of  $SN^2$  reaction  
(3) Increases the rate of  $SN^2$  reaction  
(4) Does not change the rate of  $SN^1$  and  $SN^2$  reactions
64. Which of these is the best leaving group ?  
(1) Chloride      (2) Fluoride      (3) Tosylate      (4) None
65. Of these which is least reactive ?  
(1)  $CH_2N_2$       (2)  $CH_2 = C = O$       (3)  $:CH_2$       (4)  $\cdot\dot{C}H_2$

66. By adding sodium dodecyl sulfate during the electrophoresis of proteins, it is possible to :
- (1) determine a proteins isoelectric point
  - (2) determine an enzymes specific activity
  - (3) preserve a proteins native structure
  - (4) determine the amino acid composition
67. The triplet carbene with cis-alkene gives :
- (1) cis-product
  - (2) trans-product
  - (3) both cis and trans products
  - (4) no product
68. DNFB is used to identify the N-terminal amino acid of peptides. What is this reagent called ?
- (1) Van-Slyke Reagent
  - (2) Sorenson Reagent
  - (3) Sanger's Reagent
  - (4) None of these
69. Internal reference for F-19 is :
- (1)  $NaF$
  - (2)  $CF_4$
  - (3)  $CFCl_3$
  - (4)  $NH_4F$
70. What is the internal reference for N- 15 ?
- (1) Liq.  $NH_3$
  - (2)  $NH_4OH$
  - (3)  $NH_4Cl$
  - (4)  $NH_4F$
71. What is *incorrect* for  $SN^1$  reactions ?
- (1) Rearrangement is possible
  - (2) Rate is affected by solvent polarity
  - (3) The strength of the nucleophile is important in determining the rate
  - (4) The order of reactivity is  $3^\circ > 2^\circ > 1^\circ$
72. Number of orientations with respect to applied magnetic field for deuterium is :
- (1) 2
  - (2) 3
  - (3) 1
  - (4) 4
73. Aspartic acid shows :
- (1)  $pK_{a1}$
  - (2)  $pK_{a2}$
  - (3)  $pK_{a1}$  and  $pK_{a2}$
  - (4)  $pK_{a1}$ ,  $pK_{a2}$  and  $pK_{a3}$

74. Which is *incorrect* regarding grading of sugars ?  
(1) Sucrose-1      (2) Fructose-1.75      (3) Lactose-6      (4) Saccharin-3500
75. In trimethylanilinium cation, the o, m and p-protons are deshielded because of :  
(1) Resonance      (2) Inductive effect  
(3) Both of these      (4) None of these
76. The protons of the middle carbon in allyl carbanion absorb at what ppm ?  
(1) 2.46      (2) 4.75      (3) 1.5      (4) 6.28
77. Which of these enhances the absorption of Vitamin A ?  
(1) Vit. E      (2) Vit. K      (3) DMG      (4) None
78. The CH proton in isopropyl carbocation absorbs at what ppm ?  
(1) 5.06      (2) 6.28      (3) 4.75      (4) 13.50
79. What is the *correct* decreasing order of reactivity towards electrophilic aromatic substitution ?  
(1) Indole > Pyrrole > Pyridine      (2) Pyrrole > Pyridine > Indole  
(3) Pyrrole > Indole > Pyridine      (4) Indole > Pyridine > Pyrrole
80. Which is an Anti-cancer drug ?  
(1) Camptothecin      (2) Captopril      (3) Carprofen      (4) Etodolac
81. Which is a formyl anion equivalent ?  
(1) 1,4-dithiane      (2) Ethyl chloroformate  
(3) Nitromethane      (4) Acetylene
82. The CH proton in allyl carbocation absorbs at what ppm ?  
(1) 2.56      (2) 9.64      (3) 8.97      (4) 3.56
83. The carboxypeptidase enzyme contains :  
(1) Zinc (II) and hydrolyzes COO bond  
(2) Mg (II) and hydrolyzes COO bond  
(3) Zinc (II) and hydrolyzes peptide bond  
(4) Mg (II) and hydrolyzes peptide bond

84. What is *correct* about relaxation times ?  
(1)  $T_2 = T_1$       (2)  $T_2 > T_1$       (3)  $T_2 < T_1$       (4) None of these
85. CMR spectrum of camphor shows how many peaks for carbons ?  
(1) 10      (2) 9      (3) 8      (4) 7
86. Magic angle in degrees along z-direction in NMR is :  
(1) 45.1      (2) 54.7      (3) 135.3      (4) 125.3
87. Boltzmann population excess for protons at 800 MHz at 298 K is :  
(1) 16      (2) 32      (3) 48      (4) 128
88. ROH signal appears at about what ppm range in NMR ?  
(1) 0.5 – 5.0      (2) 0.1 – 8.0      (3) 0.3 – 4.0      (4) 0.3 – 10.0
89. Which aromatic band shows fine structure in UV spectrum ?  
(1) primary      (2) secondary      (3) tertiary      (4) none
90. What is *incorrect* about electron impact ionization technique ?  
(1) It always leads to the appearance of the parent peak in the mass spectrum  
(2) It is a "hard" technique  
(3) It involves more fragmentation of the parent peak in comparison to the Chemical Ionization (CI) technique  
(4) It involves a potential of 50 – 70 eV for ionization
91. Which is *correct* pair about the use of a matrix in mass spectrometry ?  
(1) ESI + MALDI      (2) FAB + MALDI  
(3) EI + MALDI      (4) CI + MALDI
92. The NH protons in coproporphyrin absorb at about what ppm ?  
(1) + 1.0      (2) (-) 2.0      (3) (+) 3.0      (4) (-) 4.0
93. The general chemical shift range for P-31 in ppm is :  
(1) 0 - 500      (2) 0 - 600      (3) 0 - 700      (4) 0 – 1000
94. The number of orientations for B-11 with respect to applied field is :  
(1) 2      (2) 3      (3) 4      (4) 5

95. Continuous wave NMR spectroscopy gives a spectrum which is :  
(1) Frequency domain (2) Time domain  
(3) Both of these (4) None of these
96. Which is a better Diels-Alder diene for reaction with maleic anhydride ?  
(1) Furan (2) Pyrrole (3) Thiophene (4) Pyridine
97. The thermal ring opening reactions of cyclobutenes are :  
(1) Conrotatory  
(2) Disrotatory  
(3) Conrotatory or disrotatory depending upon the reaction temperature  
(4) Cannot be predicted
98. Which is *not* used in treatment of arthritis ?  
(1) Glucosamine sulfate (2) Chondroitin sulfate  
(3) Methylsulfonyl methane (4) Tosylchloride
99. Hexene-1 after reaction with metachloroperbenzoic acid followed by treatment with Lithium aluminium hydride and then with water in acidic medium gives :  
(1) Hexane (2) Hexan-1-ol (3) Hexan-2-ol (4) None
100. Betaine is an intermediate in :  
(1) Wittig Reaction (2) Stobbe Reaction  
(3) Stephenson Reduction (4) MPV Reduction