for The Young of John ?

Total No. of Printed Pages: 13

(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2013

SUBJECT: Chemistry

| C / | | Sr. No |
|---------------------------------------|-------------------------------------|--------------------------------|
| Time : 1¼ Hours Roll No. (in figures) | Max. Marks : 100 (in words) | Total Questions: 100 |
| Name | Father's Name Date of Examination _ | |
| (Signature of the Candidate) | | (Signature of the Invigilator) |

CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

- 1. All questions are compulsory and carry equal marks.
- 2. The candidates must return the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
- 3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing within two hours after the test is over. No such complaint(s) will be entertained thereafter.
- 4. The candidate *must not* do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers Must Not be ticked in the question booklet.
- 5. Use only black or blue ball point pen of good quality in the OMR Answer-Sheet.
- 6. There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.
- 7. Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet, containing 100 questions (Sr. No. 1 to 100). Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.

PG-EE-2013/Chemistry/(C)

| 1. | In Rutile structure, the coordination num | nbei | of Titanium ato | ms i | s: |
|----|---|------|-------------------------|-------|------------------|
| | (1) Six (2) Four | (3) | Two | (4) | Eight |
| 2. | Which of the following metal ion pairs h | nave | similar ionic rac | lii? | |
| | (1) Ti^{4+} and Zr^{4+} | (2) | V^{5+} and Nb^{5+} | | |
| | (3) Cr^{3+} and Mn^{3+} | (4) | Zr^{4+} and Hf^{4+} | | over some f |
| 3. | Which of the following solid will behave | e as | p-type semicond | lucto | or? |
| | (1) NaCl (2) ZnS | (3) | FeS | (4) | AgCl |
| 4. | Which metal has highest cohesion energ | y? | | | |
| | (1) Cobalt (2) Nickel | (3) | Copper | (4) | Zinc |
| 5. | The aqueous solution of which metal ion | n wi | ll be colourless? | | |
| | (1) Ti^{3+} (2) Cr^{3+} | (3) | Cu ⁺ | (4) | Cu^{2+} |
| 6. | Which of the following is a Lanthanide | elen | nent? | | |
| | (1) Francium (2) Europium | (3) | Tungsten | (4) | Polonium |
| 7. | In the reaction $HClO_4 + HF \rightleftharpoons H_2F^+ +$ | ClO | the base is: | | |
| | (1) HClO ₄ (2) HF | (3) | H_2F^+ | (4) | ClO ₄ |
| 8. | Which of the following will behave as a | Lev | vis acid? | | |
| | (1) NH_3 (2) NH_4^+ | (3) | BF ₃ | (4) | CH ₄ |
| 9. | If you titrate an aqueous solution of bor | ax v | vith HCl, indicate | or us | sed will be: |
| | (1) Phenolphthalein | (2) | Methyl orange | | |
| | (3) Methyl red | (4) | Eriochrome bla | ck T | |
| 0. | As per HSAB concept, the hardest acid | will | be: | | and y |
| | (1) Fe^{3+} (2) Zn^{2+} | (3) | Ag^+ | (4) | Hg ²⁺ |
| 1. | Which is a local anaesthetic? | | | | 10270000 |
| | (1) Cocaine (2) Quinine | (3) | Morphine | (4) | None |

| 12 | . Which enhances the absorption of Vit | amin A ? | |
|-------|--|--|------|
| | (1) Vit. K (2) Vit. C | (3) DMG (4) None | |
| 13. | By which of the following reaction, ac | etophenone can be converted to phenol ? | |
| | (1) m-CPBA followed by base catalyz | | |
| | (2) Conc. HNO ₃ | | |
| | (3) Iodine and NaOH | | |
| | (4) Singlet oxygen followed by hydro | lysis | |
| 14. | Diazomethane with acetylene gives: | Agent Comment of the American Beauty States | |
| | (1) Pyrazole (2) Pyrazoline | (3) Piperidine (4) Pyrimidine | |
| 15. | Cinnamoyl alcohol with lead tetraaceta | ate gives : | |
| | (1) Cinnamic acid | (2) Cinnamoyl acetate | |
| | (3) Cinnamaldehyde | (4) Acetophenone | |
| 16. | Betaine is an intermediate in : | | |
| | (1) Wittig reaction | (2) Stobbe reaction | |
| | (3) Stephenson reduction | (4) MPV reduction | |
| 17. | If the migrating group in Beckman rear | rrangement is chiral, then: | |
| | (1) Its configuration will change | | |
| | (2) Its configuration will be retained | | |
| | (3) Both | | |
| | (4) None | | |
| 18. | Which reduces only the carbonyl grobond and ester functional groups? | up in the presence of nitro, carboxyl, dou | ıble |
| | | (3) NaBH ₄ (4) H ₂ /Ni | |
| PG-EE | -2013/Chemistry/(C) | | |

| 19. Which is the correct decreasing substitution? | order of reactivity towards electrophilic aromatic |
|---|--|
| (1) Indole > Pyrrole > Pyridine | (2) Pyrrole > Pyridine > Indole |
| (3) Pyrrole > Indole > Pyridine | (4) Indole > Pyridine > Pyrrole |
| 20. OH signal of alcohol appears at wh | at ppm range? |
| (1) 0.5 – 5.0 (2) 0.1 – 8.0 | (3) 0.3 – 4.0 (4) 0.3 – 10.0 |
| 21. C = C frequency in Oct-4-ene appear | rs at: |
| (1) 1680-1600 cm ⁻¹ (very weak) | confolion confolion |
| (2) 1680-1600 cm ⁻¹ (strong) | with acety |
| (3) 1680-1600 cm ⁻¹ (m) | (2) 7 |
| (4) No peak in this region of 1680-16 | 600 cm ⁻¹ |
| 22. I for C-13 is: | , |
| (1) 1 (2) 1/2 | (3) 3/2 (4) 2 |
| 23. I for P-31 is: | alsibratures in a second second |
| (1) 1 (2) 1/2 | (3) 3/2 (4) 3 |
| 24. What is the right order of coupling co | onstants? |
| (1) $J^1 > J^2 > J^3$ (2) $J^3 > J^2 > J^1$ | (3) $J^1 = J^2 = J^3$ (4) None of these |
| 25. Which aromatic band shows fine structure | cture? |
| (1) Primary (2) Secondary | (3) Tertiary (4) None |
| 26. Which is a better Diels Alder Diene for | |
| (1) Furan (2) Pyrrole | (3) Thiophene (4) Pyridine |
| 27. Which is a strong base? | |
| (1) Aniline | (2) Cyclohexylamine |
| (3) Pyrrole | (4) Quinoline |
| PG-EE-2013/Chemistry/(C) | |
| | P. T. O. |

(1)
$$CH_3 - CH_2 > OH_2 > CH = C > OH$$

(2)
$$CH \equiv C > NH_2 > CH \equiv C > OH$$

(3)
$$OH > NH_2 > CH \equiv C > CH_3 - CH_2$$

(4).
$$\stackrel{\Theta}{NH_2} > CH \equiv \stackrel{\Theta}{C} > \stackrel{\Theta}{OH} > CH_3 - \stackrel{\Theta}{CH_2}$$

Which gives single mononitroderivative?

- (1) Naphthalene (2) O-xylene
- (3) Ethylbenzene (4) p-xylene

Which one is most effective in an SN^2 displacement on methyl bromide?

- (1) C₂H₅ $\overset{\Theta}{O}$
- (2) HO
- (3) $C_6H_5\overset{\Theta}{O}$ (4) $CH_3CO\overset{\Theta}{O}$

The force constant of a diatomic S.H.O. can be calculated by employing relation:

 $(1) \quad k = 4\pi^2 c^2 \left(\overline{v}^2\right) \mu$

(2) $k = 4\pi^2 c(\overline{v}^2)\mu$

- (3) $k = 4\pi^2 c(\overline{v}) \mu^2$
- (4) $k = 4\pi^2 \,\mu \,c$

where all the symbols have their usual meaning.

Zero point energy for diatomic molecule possessing harmonic motion is : 32.

- (1) zero
- (3) $\frac{1}{2}hv$ (4) $\frac{1}{3}hv$

The power output of a laser in which 2.0 J pulse can be delivered in one nanosecond

- (1) 2.0 GW
- (2) 20.0 GW
- (3) 0.20 GW
- (4) None of these

For Arhenius equation, $A = e^{-E_a/RT}$, if $T \to \infty$, then value of E_a will be:

- (1) positive
- (2) negative
- (3) zero
- (4) equal to A

The molarity of pure water is:

- (1) 50
- (2) 18
- (3) 100
- (4) 55.6

PG-EE-2013/Chemistry/(C)

P. T. O.

| 36 | The degeneracy of the rotational ener is: | gy level with J | = 4 for a heterodiatomic molecul | e |
|-------|---|--------------------|--|---|
| | (1) 4 (2) 7 | (3) 9 | (4) 8 | |
| 37 | . Mean free path of a gas molecule is: | €. | ED < LHX | |
| | (1) inversely proportional to pressure | e | S. Hoer and the same | |
| | (2) directly proportional to pressure | ¥- | | |
| | (3) independent of pressure | | n case of college at the | |
| | (4) independent of temperature | | donale · · · · · · · · · · · · · · · · · · · | |
| 38. | In B.E.T. equation one of the following | g statement is no | ot true. Select the one: | |
| | (1) It considers the multi layer adsorp | otion | tride tech in E. (4) | |
| | (2) It doesn't use the concept of satura | ation of vapour | pressure | |
| | (3) It is not valid for porous adsorben | t | | |
| | (4) It uses the concept of latent heat o | f condensation | | |
| 39. | No diffraction would result, if: | | tenno pilko neli Man solt. 84. | |
| | (1) $\lambda < 2d$ (2) $\lambda \approx 2d$ | (3) λ < < d | (4) $\lambda > 2d$ | |
| 40. | 11.2×10^3 m ³ of a gas at STP requires 1 C_v for the gas is : | 04.6 J to raise it | s temperature by 10 degree. The | |
| | (1) 20.92 J deg ⁻¹ mole ⁻¹ | (2) 10.46 J de | eg ⁻¹ mole ⁻¹ | |
| | (3) $9.4 J \text{deg}^{-1} \text{mole}^{-1}$ | (4) zero | | |
| 41. | How many peaks are observed in UV- | visible absorption | on spectra of $\left[Ni(H_2O)_6^2\right]^{2+}$? | |
| | (1) One (2) Two | (3) Three | (4) Four | |
| 42. | Write the Ground Term of Cr^{3+} : | | | |
| | (1) 6_S (2) 4_F | (3) 2 _D | (4) 3 _P | |
| PG-EE | -2013/Chemistry/(C) | | P. T. O | |

| 43. | Predict the Point | Group in | $Fe(CO)_5$: |
|-----|-------------------|----------|--------------|
|-----|-------------------|----------|--------------|

- (1) Oh
- (2) C3V
- (3) C_{2V}
- (4) D_{3h}

Nitrogenase enzyme consists of:

- (1) Co
- (2) Se
- (3) Mo, Fe
- (4) Mg

45. Vitamin
$$B_{12}$$
 consists of:

- (1) Fe
- (2) Co 🝜 -
- (3) Mn (4) V

$$\int_{92}^{235} U + \int_{0}^{1} n \to_{56}^{141} Ba + \int_{36}^{92} Kr + \dots$$

- $(1) \ 2 \ \frac{1}{0}n$
- $(2) \frac{1}{1}H$
- (3) ${}_{1}^{2}H$
- (4) ⁴₂He

Bhopal Tragedy which killed thousands of people, was due to air pollution of:

(1) CO

(2) SO₂

(3) Nitrogen oxides

(4) Methyl Isocyanate

- (1) $\hat{L}_x = i\hbar \left[x \cdot \frac{\partial}{\partial x} z \cdot \frac{\partial}{\partial z} \right]$ (2) $-i\hbar \left[y \cdot \frac{\partial}{\partial z} z \cdot \frac{\partial}{\partial y} \right]$
- (3) $\hat{L}_x = i\hbar \left[y \cdot \frac{\partial}{\partial z} z \cdot \frac{\partial}{\partial u} \right]$
- (4) $-i\hbar \left[x \cdot \frac{\partial}{\partial z} z \cdot \frac{\partial}{\partial x}\right]$

49. Operators
$$\hat{A}$$
 and \hat{B} are said to be commutative, if :

(1) $\hat{A} - \hat{B} = 0$

(2) $\hat{A} + \hat{B} = 0$

(3) $\hat{A} \hat{B} - \hat{B} \hat{A} = 0$

(4) $\hat{A} \hat{B} + \hat{B} \hat{A} = 0$

- (1) $\frac{\sqrt{2}}{a}\sin\frac{n\pi x}{a}$ (2) $\sqrt{\frac{2}{a}}\frac{n\pi x}{a}$ (3) $\sqrt{\frac{2}{a}}\sin\frac{\pi x}{a}$ (4) $\sqrt{\frac{2}{a}}\sin\frac{n\pi x}{a}$

| 51. | What is the decreas | sing order of chemic | al shi | fts for protons a | mor | ng these ? | |
|-------|-------------------------------------|--|--------|-----------------------------|--------|-------------------|----------|
| | | nnes > Alkenes | | Alkanes > Alke | | | |
| | (3) Alkynes > Alke | | | Alkenes > Alky | nes | > Alkanes | |
| 52. | The singlet at about which protons? | t 4.0 ppm in the pro | oton l | NMR spectrum | of m | nethylacetate is | due to |
| | (1) Methyl | | (2) | Methoxy | | | |
| | (3) Methyl and Mo | ethoxy | (4) | None of these | | | |
| 53. | Which is not an ant | ti-cancer drug? | | | | verse and made | |
| | (1) Vincristine | 9.1 | (2) | Cyclophosphai | mide | | |
| | (3) Doxorubicin | 31 | (4) | Gabapentin | | | |
| 54. | | ction with metachlor hydride and then w | | | | | nt with |
| | (1) Hexane | (2) Hexan-1-ol | (3) | Hexan-2-ol | (4) | None | |
| 55. | Write the symbol o | f atomic orbital if n : | = 3, 1 | = 2 and $m = -2$, | -1, | 0, + 1, + 2: | |
| | (1) 2s | (2) 3s | (3) | 3p | (4) | 3 <i>d</i> | |
| 56. | An element with at | omic number 72 bel | ongs | to: | | | |
| | (1) s-block | (2) p-block | (3) | d-block | (4) | f-block | |
| 57. | Which of the follow | ving metals has lowe | est io | nization potenti | al? | | |
| | (1) Lithium | (2) Sodium | (3) | Berylium | (4) | Magnesium | |
| 58. | Which cation has h | ighest polarizing po | wer ' | ? | | | |
| | (1) Na ⁺ | (2) Mg^{2+} | (3) | K ⁺ | (4) | Al^{3+} | |
| 59. | How many lone pa | irs of electrons are p | oresei | nt in ICl ₂ ion? | | | |
| | (1) Zero | (2) One | (3) | Two | (4) | Three | |
| 60. | Which of the follow | ving molecules/ions | has | smallest O – O l | ond | ? | |
| | (1) O ₂ | (2) O ₂ ⁺ , | (3) | O ₂ | (4) | O_2^{2-} | |
| 61. | The Boyle tempera | ture is that at which | the s | econd virial coe | fficie | ent of real gas i | s: |
| | (1) zero | (2) one | (3) | four | (4) | one and half | |
| PG-EE | E-2013/Chemistry/(C |) | | | | | P. T. O. |

| 62. | The fugacity function is defined as: | desirant harabas various site algorithms to the se |
|-------|---|--|
| | (1) $\underset{P\to 0}{limit} \frac{p}{f} = 1$ (2) $\underset{P\to 0}{limit} \frac{f}{p} = 1$ | (3) $\underset{f\to 0}{limit} \frac{p}{f} = 1$ (4) $\underset{P\to 0}{limit} \frac{p}{f} = 0$ |
| 63. | Choose the correct relation: | |
| | | (2) $\left(\frac{\partial A}{\partial T}\right)_V = \left(\frac{\partial G}{\partial T}\right)_P$ |
| | (3) $\left(\frac{\partial T}{\partial S}\right)_P = \left(\frac{\partial V}{\partial S}\right)_P$ | (4) $\left(\frac{\partial S}{\partial P}\right)_T = -\left(\frac{\partial T}{\partial V}\right)_P$ |
| 64. | For the combustion of one mole of CF | I ₃ COOH(l) at 298 K, Δn is : |
| | (1) 1 (2) -1 | (3) zero (4) -1/2 |
| day. | In the limit $T \rightarrow 0$, for a crystal: | |
| | where C_p is the heat capacity at const | (3) $S_T = C_p / 2$ (4) $S_T = C_p / 3$ |
| cc | | |
| 66. | | Entire Transfer of the party of the same o |
| | (1) 0.375 (2) 0.400 | (3) zero (4) 0.512 |
| 67. | The Joule-Thomson expansion of an ic | leal gas is : |
| | (1) Adiabatic process | (2) an isentropic process |
| | (3) an isenthalpic process | (4) an isothermal process |
| 68. | The spacing between 123 planes in a $b = 100$ pm and $c = 150$ pm is: | an orthorhombic unit cells having $a = 50$ pm |
| | (1) 2.9 pm (2) 29 pm | (3) 9.2 pm (4) 92 pm |
| 69. | The cell potential is a: | |
| | (1) Colligative property | (2) Thermodynamic property |
| | (3) Intensive property | (4) Extensive property |
| 70. | The solubility of silver chloride in wat product will be: | er at 298.15 K is $0.00179 \text{ g litre}^{-1}$. The solubility |
| | (1) $156 \times 10^{-10} \mathrm{mol}^2 \mathrm{dm}^{-6}$ | (2) $1.56 \times 10^{-9} \text{ mol}^2 \text{dm}^{-6}$ |
| | (3) $15.6 \times 10^{-12} \mathrm{mol}^2 \mathrm{dm}^{-6}$ | (4) $1.56 \times 10^{-10} \mathrm{mol}^2 \mathrm{dm}^{-6}$ |
| PG-EE | -2013/Chemistry/(C) | Sittle (b) Site (1) Site (1) |

| 71. | In the lead acid bat | tery during charging | , the | cathode reaction | n is: | |
|-----|---|---|--------|-----------------------------------|------------------|--|
| | (1) reduction of Pl | o ⁺² to Pb | (2) | formation of Pl | bSO ₄ | |
| | (3) formation of Pi | bO ₂ | (4) | None of these | | |
| 72. | When a radioactive | e element loses one 'o | a' and | d two 'β' particle | es, it y | yields: |
| | (1) Isobar | (2) Isomer | (3) | Isotope | (4) | Allotrope |
| 73. | 50 ml of 0.1 NaOH | are added to 49 ml o | | | the re | esulting solution is |
| | (1) 12 | (2) 11 | (3) | 10 | (4) | 9 |
| 74. | The heat of reaction | n is independent of: | | | | |
| | (1) Pressure | 7. | (2) | Temperature | | |
| | (3) Physical state | | (4) | The path by wl | hich 1 | product is formed |
| 75. | Which of the follow | wing will show ESR s | spect | ra? | | |
| | (1) C_6H_6 | (2) CH ₃ | (3) | CH ₄ | (4) | H_2 |
| 76. | What is the freque | ncy of radiation poss | sessir | ng wave length | 400 n | m? |
| | (1) $7.5 \times 10^{-14} \mathrm{S}^{-1}$ | (2) $7.5 \times 10^{14} \text{ S}^{-1}$ | (3) | $7.5 \times 10^{9} \text{S}^{-1}$ | (4) | $7.5 \times 10^{-13} \text{ S}^{-1}$ |
| 77. | In aerosol, the disp | persion medium is: | | Rosensons Edit | | Continue to Contin |
| | (1) Gas | | (3) | Liquid | (4) | Mixture of all |
| 78. | The polymers cons | sist of coil like polym | ner cl | nain are: | | Personal III |
| | (1) Thermoplasts | (2) Elastomers | (3) | Thermosets | (4) | None of these |
| 79. | Which of the follow | wing is a state functi | on? | | | |
| | (1) E – PV | (2) E + PV | (3) | Q/W | (4) | Q - W |
| 80. | The ilkovic equati | on for diffusion curr | ent is | s expressed as: | | proceedings (g) |
| | $(1) \overrightarrow{I}_d = 607n DC$ | $2m^{2/3}t^{1/6}$ | | $\overrightarrow{I}_d = 607n D^1$ | | |
| | $(3) \overrightarrow{I}_d = 607nCD$ | $n^{1/2} m^{2/3} t^{1/6}$ | (4) | $\overrightarrow{I}_d = 607n D^1$ | $^{/2}C^{1}$ | $^{/2} m^{1/3} t^{1/6}$ |

| 81 | . Which reacts fastest with N-bromos | uccinimide (NBS) ? |
|--------|---|--|
| | (1) Toluene (2) Methane | (3) Pyridine (4) Benzene |
| 82 | . When vinyl cyanide reacts with ethy | valled a presence of a base, what is formed |
| | $(1) CH_2 = CH - OH$ | (2) $C_2H_5O - CH_2 - CH_2CN$ |
| | (3) CH ₃ CH ₂ OH | (4) $C_2H_5 - O - C_2H_5$ |
| 83. | Which is the best leaving group? | |
| | (1) Chloride (2) Fluoride | (3) Tosylate (4) None |
| 84. | With cis-alkenes, the triplet carbenes | give: |
| | (1) cis-product | (2) trans-product |
| | (3) no product | (4) both cis and trans products |
| 85. | DNFB is used to identify N-terminal | amino acid of peptides. The reagent is called: |
| | (1) Van-Slyke reagent | (2) Sorenson reagent |
| | (3) Sanger's reagent | (4) Stephens reagent |
| 86. | Continuous wave NMR spectroscopy | involves: |
| | (1) sequential detection of resonances | s of nuclei |
| | (2) simultaneous detection of all reso | nances of nuclei |
| | (3) sometimes sequential and sometimes | mes simultaneous detection of nuclei |
| | (4) None | |
| 87. | The addition of Br_2 to methyl acetyler | ne to give trans-1, 2-dibromopropene is a : |
| | (1) Stereoselective reaction | |
| | (2) Stereospecific reaction | |
| | (3) Stereoselective and Stereospecific | reaction |
| | (4) None | |
| PG-EE- | -2013/Chemistry/(C) | and the second s |

P. T. O.

C

PG-EE-2013/Chemistry/(C)

Predict the type of isomerism in $[Co(NH_3)_6][Cr(CN)_6]$ and $[Cr(NH_3)_6][Co(CN)_6]$:

- (1) Linkage Isomerism (2) Coordination Isomerism
- (3) Stereoisomerism (4) Coordination position Isomerism

Which of the following complex ions will not be square planar in structure?

- (1) $\left[Co(CN)_4\right]^{2-}$ (2) $\left[Ni(CN)_4^{*}\right]^{2-}$ (3) $\left[Cu(NH_3)_4\right]^{2+}$ (4) $Ni(CO)_4$